

Boundedness of maximal operators and Sobolev's theorem for non-homogeneous central Morrey spaces of variable exponent

Yoshihiro Mizuta, Takao Ohno and Tetsu Shimomura

Abstract

Our aim in this paper is to deal with the boundedness of the Hardy-Littlewood maximal operator in non-homogeneous central Morrey spaces of variable exponent. Further, we give Sobolev's inequality and Trudinger's exponential integrability for generalized Riesz potentials.

1 Introduction

Let \mathbf{R}^N be the Euclidean space. In [4], Beurling introduced the space $B^p(\mathbf{R}^N)$ to extend Wiener's ideas [21, 22] which describes the behavior of functions at infinity. Feichtinger [8] gave an equivalent norm on $B^p(\mathbf{R}^N)$, which is a special case of norms in Herz spaces $K_p^{\alpha,r}(\mathbf{R}^N)$ introduced by Herz [12]. Precisely speaking, $B^p(\mathbf{R}^N) = K_p^{-N/p,\infty}(\mathbf{R}^N)$ (see also [11]). In [10], García-Cuerva studied the boundedness of the maximal operator on the space $B^p(\mathbf{R}^N)$. As an extension of the space $B^p(\mathbf{R}^N)$, García-Cuerva and Herrero [11] introduced the central Morrey spaces $B^{p,\nu}(\mathbf{R}^N)$ (see also [3]). Alvarez, Guzmán-Partida and Lakey [3] obtained the boundedness of a class of singular integrals operators on the central Morrey spaces (see also Komori [13]), which are more singular than Calderón-Zygmund operators and include pseudo-differential operators.

Variable exponent Lebesgue spaces and Sobolev spaces were introduced to discuss nonlinear partial differential equations with non-standard growth condition. Our first aim in this paper is to introduce the non-homogeneous central Morrey spaces of variable exponent, and study the boundedness of the Hardy-Littlewood maximal operator (see Theorem 3.1), in a way different from Almeida and Drihem [2].

In classical Lebesgue spaces, we know Sobolev's inequality :

$$\|I_\alpha f\|_{L^{p^\#}(\mathbf{R}^N)} \leq C \|f\|_{L^p(\mathbf{R}^N)}$$

2000 Mathematics Subject Classification : Primary 31B15, 46E35

Key words and phrases : Maximal operator, non-homogeneous central Morrey spaces of variable exponent, Riesz potentials, Sobolev's theorem, Sobolev's inequality, Trudinger's exponential integrability

for $f \in L^p(\mathbf{R}^N)$, $0 < \alpha < N$ and $1 < p < N/\alpha$, where I_α is the Riesz kernel of order α and $1/p^\sharp = 1/p - \alpha/N$ (see, e.g. [1, Theorem 3.1.4]). This result was extended to the central Morrey spaces by Fu, Lin and Lu [9, Proposition 1.1] (see also Matsuoka and Nakai [15]).

To obtain general results, for $0 < \alpha < N$ and an integer k , we define the generalized Riesz potential $I_{\alpha,k}f$ of order α of a locally integrable function f on \mathbf{R}^N by

$$I_{\alpha,k}f(x) = \int_{\mathbf{R}^N \setminus B(0,1)} \left\{ I_\alpha(x-y) - \sum_{\{\mu: |\mu| \leq k-1\}} \frac{x^\mu}{\mu!} (D^\mu I_\alpha)(-y) \right\} f(y) dy,$$

where $I_\alpha(x) = |x|^{\alpha-n}$ (see [16, 17]). Remark here that

$$I_{\alpha,k}f(x) = \int_{\mathbf{R}^N \setminus B(0,1)} I_\alpha(x-y) f(y) dy$$

when $k \leq 0$.

In Section 4, when $p^+ < N/\alpha$ (see Section 2 for the definition of p^+), we shall give Sobolev's inequality for $I_{\alpha,k}f$ with functions in the non-homogeneous central Morrey spaces of variable exponent (see Theorem 4.5); for related result, we refer the reader to Fu, Lin and Lu [9, Theorem 2.1].

In the last section, when $p = N/\alpha$, we treat Trudinger's exponential integrability for $I_{\alpha,k}f$ (see Theorem 5.1).

2 Preliminaries

Consider a function $p(\cdot)$ on \mathbf{R}^N such that

$$(P1) \quad 1 < p^- := \inf_{x \in \mathbf{R}^N} p(x) \leq \sup_{x \in \mathbf{R}^N} p(x) =: p^+ < \infty;$$

(P2) $p(\cdot)$ is log-Hölder continuous, namely

$$|p(x) - p(y)| \leq \frac{c_p}{\log(e + 1/|x - y|)} \quad \text{for } x, y \in \mathbf{R}^N$$

with a constant $c_p \geq 0$;

(P3) $p(\cdot)$ is log-Hölder continuous at ∞ , namely

$$|p(x) - p(\infty)| \leq \frac{c_\infty}{\log(e + |x|)} \quad \text{whenever } |x| > 0$$

with constants $p(\infty) > 1$ and $c_\infty \geq 0$;

$p(\cdot)$ is referred to as a variable exponent.

For $\nu \geq 0$, we denote by $\mathcal{B}^{p(\cdot), \nu}(\mathbf{R}^N)$ the class of locally integrable functions f on \mathbf{R}^N satisfying

$$\|f\|_{\mathcal{B}^{p(\cdot), \nu}(\mathbf{R}^N)} = \sup_{R \geq 1} R^{-\nu/p(\infty)} \|f\|_{L^{p(\cdot)}(B(0,R))} < \infty,$$

where

$$\|f\|_{L^{p(\cdot)}(B(0,R))} = \inf \left\{ \lambda > 0 : \int_{B(0,R)} \left(\frac{|f(y)|}{\lambda} \right)^{p(y)} dy \leq 1 \right\}.$$

The space $\mathcal{B}^{p(\cdot),\nu}(\mathbf{R}^N)$ is referred to as a non-homogeneous central Morrey spaces of variable exponent. If $p(\cdot)$ is a constant and $\nu = N$, then $\mathcal{B}^{p(\cdot),\nu}(\mathbf{R}^N) = B^p(\mathbf{R}^N)$.

Throughout this paper, let C denote various constants independent of the variables in question. The symbol $g \sim h$ means that $C^{-1}h \leq g \leq Ch$ for some constant $C > 0$.

LEMMA 2.1. *Set*

$$\|f\|_{\tilde{\mathcal{B}}^{p(\cdot),\nu}(\mathbf{R}^N)} = \inf \left\{ \lambda > 0 : \sup_{R \geq 1} R^{-\nu} \int_{B(0,R)} \left(\frac{|f(y)|}{\lambda} \right)^{p(y)} dy \leq 1 \right\}.$$

Then

$$\|f\|_{\mathcal{B}^{p(\cdot),\nu}(\mathbf{R}^N)} \sim \|f\|_{\tilde{\mathcal{B}}^{p(\cdot),\nu}(\mathbf{R}^N)}$$

for all $f \in L^1_{loc}(\mathbf{R}^N)$.

Proof. We may assume that $\nu > 0$. First we find a constant $C > 0$ such that

$$\|f\|_{\mathcal{B}^{p(\cdot),\nu}(\mathbf{R}^N)} \leq C \|f\|_{\tilde{\mathcal{B}}^{p(\cdot),\nu}(\mathbf{R}^N)}$$

for all $f \in L^1_{loc}(\mathbf{R}^N)$. Let f be a nonnegative function on \mathbf{R}^N with $\|f\|_{\tilde{\mathcal{B}}^{p(\cdot),\nu}(\mathbf{R}^N)} \leq 1$. Then note that

$$R^{-\nu} \int_{B(0,R)} f(y)^{p(y)} dy \leq 1$$

for all $R \geq 1$. To end the proof, it is sufficient to find a constant $C > 0$ such that

$$\int_{B(0,R) \setminus B(0,1)} (R^{-\nu/p(\infty)} f(y))^{p(y)} dy \leq C$$

for all $R \geq 1$. For this purpose, let $R \geq 1$ and take an integer $j_0 \geq 1$ such that $2^{-j_0} R \leq 1 < 2^{-j_0+1} R$. We have

$$\begin{aligned} & \int_{B(0,R) \setminus B(0,1)} (R^{-\nu/p(\infty)} f(y))^{p(y)} dy \\ & \leq \sum_{j=0}^{j_0} \int_{B(0,2^{-j+1}R) \setminus B(0,2^{-j}R)} (R^{-\nu/p(\infty)} f(y))^{p(y)} dy \\ & \leq \sum_{j=0}^{j_0} (2^{-j})^{\nu/p(\infty)} \int_{B(0,2^{-j+1}R) \setminus B(0,2^{-j}R)} \{(2^{-j}R)^{-\nu/p(\infty)} f(y)\}^{p(y)} dy \\ & \leq \sum_{j=0}^{j_0} (2^{-j})^{\nu/p(\infty)} (2^{-j}R)^\nu \int_{B(0,2^{-j+1}R)} f(y)^{p(y)} dy \\ & \leq C \sum_{j=0}^{j_0} (2^{-j})^{\nu/p(\infty)} \leq C \end{aligned}$$

since $|y|^{-p(y)} \leq C|y|^{-p(\infty)}$ for $y \in B(0, 2^{-j+1}R) \setminus B(0, 2^{-j}R)$ and $0 \leq j \leq j_0$ by (P3).

Next we prove the converse inequality. Then it is sufficient to find a constant $C > 0$ such that

$$R^{-\nu} \int_{B(0,R) \setminus B(0,1)} f(y)^{p(y)} dy \leq C$$

for all $R \geq 1$ and $f \geq 0$ on \mathbf{R}^N with

$$\sup_{R>1} \int_{B(0,R)} (R^{-\nu/p(\infty)} f(y))^{p(y)} dy \leq 1.$$

For this purpose, let $R > 1$ and take an integer $j_0 \geq 1$ such that $2^{-j_0}R \leq 1 < 2^{-j_0+1}R$ as before. We find

$$\begin{aligned} & \int_{B(0,R) \setminus B(0,1)} (R^{-\nu/p(y)} f(y))^{p(y)} dy \\ & \leq \sum_{j=0}^{j_0} (2^{-j})^\nu \int_{B(0,2^{-j+1}R) \setminus B(0,2^{-j}R)} \{(2^{-j}R)^{-\nu/p(y)} f(y)\}^{p(y)} dy \\ & \leq \sum_{j=0}^{j_0} (2^{-j})^\nu \int_{B(0,2^{-j+1}R)} \{(2^{-j}R)^{-\nu/p(\infty)} f(y)\}^{p(y)} dy \\ & \leq C \sum_{j=0}^{j_0} (2^{-j})^\nu \leq C \end{aligned}$$

since $|y|^{-1/p(y)} \leq C|y|^{-1/p(\infty)}$ for $y \in B(0, 2^{-j+1}R) \setminus B(0, 2^{-j}R)$ and $0 \leq j \leq j_0$ by (P3). Thus the proof is completed. \square

3 Boundedness of maximal operators

For a locally integrable function f on \mathbf{R}^N , the Hardy-Littlewood maximal function Mf is defined by

$$Mf(x) = \sup_{r>0} \frac{1}{|B(x,r)|} \int_{B(x,r)} |f(y)| dy,$$

where $B(x,r)$ is the ball in \mathbf{R}^N with center x and of radius $r > 0$, and $|B(x,r)|$ denotes its Lebesgue measure. The mapping $f \mapsto Mf$ is called the maximal operator.

The maximal operator is a classical tool in harmonic analysis and studying Sobolev functions and partial differential equations, and it plays a central role in the study of differentiation, singular integrals, smoothness of functions and so on (see [5, 13, 14, 20], etc.).

It is well known that the maximal operator is bounded in the Lebesgue space $L^p(\mathbf{R}^N)$ when $p > 1$ (see [20]). We present the boundedness of maximal operator in the central Morrey spaces of variable exponent.

THEOREM 3.1. *Let $0 \leq \nu \leq N$. Then the maximal operator $f \rightarrow Mf$ is bounded from $\mathcal{B}^{p(\cdot),\nu}(\mathbf{R}^N)$ to $\mathcal{B}^{p(\cdot),\nu}(\mathbf{R}^N)$, that is,*

$$\|Mf\|_{\mathcal{B}^{p(\cdot),\nu}(\mathbf{R}^N)} \leq C\|f\|_{\mathcal{B}^{p(\cdot),\nu}(\mathbf{R}^N)} \quad \text{for all } f \in \mathcal{B}^{p(\cdot),\nu}(\mathbf{R}^N).$$

When $0 \leq \nu < N$, this theorem is essentially proved by Almeida and Drihem [2, Corollary 4.7]. But, for the readers' convenience, we give a proof of Theorem 3.1 different from [2].

Before doing this, we prepare the following results.

LEMMA 3.2 ([7, Corollary 4.5.9]). *For all $R \geq 1$,*

$$\|1\|_{L^{p(\cdot)}(B(0,R))} \sim R^{N/p(\infty)},$$

that is, $1 \in \mathcal{B}^{p(\cdot),N}(\mathbf{R}^N)$.

LEMMA 3.3. *There exists a constant $C > 0$ such that*

$$\frac{1}{|B(0,R)|} \int_{B(0,R) \setminus B(0,R/2)} f(y) dy \leq CR^{-(N-\nu)/p(\infty)}$$

for all $R \geq 1$ and $f \geq 0$ such that $\|f\|_{\mathcal{B}^{p(\cdot),\nu}(\mathbf{R}^N)} \leq 1$.

Proof. Let f be a nonnegative function on \mathbf{R}^N such that $\|f\|_{\mathcal{B}^{p(\cdot),\nu}(\mathbf{R}^N)} \leq 1$. Then we see from Lemma 2.1 that

$$R^{-\nu} \int_{B(0,R) \setminus B(0,R/2)} f(y)^{p(y)} dy \leq C$$

for all $R \geq 1$. Hence we find by (P3)

$$\begin{aligned} & \frac{1}{|B(0,R)|} \int_{B(0,R) \setminus B(0,R/2)} f(y) dy \\ & \leq R^{-(N-\nu)/p(\infty)} + \frac{1}{|B(0,R)|} \int_{B(0,R) \setminus B(0,R/2)} f(y) \left(\frac{f(y)}{R^{-(N-\nu)/p(\infty)}} \right)^{p(y)-1} dy \\ & \leq R^{-(N-\nu)/p(\infty)} + CR^{(N-\nu)(p(\infty)-1)/p(\infty)} \frac{1}{|B(0,R)|} \int_{B(0,R) \setminus B(0,R/2)} f(y)^{p(y)} dy \\ & \leq CR^{-(N-\nu)/p(\infty)} \end{aligned}$$

for all $R \geq 1$, as required. □

We denote by χ_E the characteristic function of E .

LEMMA 3.4. *Let $0 \leq \nu \leq N$. Then there exists a constant $C > 0$ such that*

$$M(f\chi_{\mathbf{R}^N \setminus B(0,2R)})(x) \leq CR^{-(N-\nu)/p(\infty)}$$

for all $x \in B(0,R)$ with $R \geq 1$ and $f \geq 0$ with $\|f\|_{\mathcal{B}^{p(\cdot),\nu}(\mathbf{R}^N)} \leq 1$.

Proof. Let f be a nonnegative function on \mathbf{R}^N such that $\|f\|_{\mathcal{B}^{p(\cdot),\nu}(\mathbf{R}^N)} \leq 1$. Let $R \geq 1$ and $x \in B(0, R)$. We have by Lemma 3.3

$$\begin{aligned}
M(f\chi_{\mathbf{R}^N \setminus B(0,2R)})(x) &= \sup_{r>R} \frac{1}{|B(x,r)|} \int_{B(x,r) \setminus B(0,2R)} f(y) dy \\
&\leq \sup_{r>R} \frac{1}{|B(0,r)|} \sum_{\{j \geq 1: 2^j R < 2r\}} \int_{B(0,2^{j+1}R) \setminus B(0,2^j R)} f(y) dy \\
&\leq C \sup_{r>R} \frac{1}{|B(0,r)|} \sum_{\{j \geq 1: 2^j R < 2r\}} (2^{j+1}R)^{N-(N-\nu)/p(\infty)} \\
&\leq C \sup_{r>R} \frac{1}{|B(0,r)|} r^{N-(N-\nu)/p(\infty)} \\
&\leq CR^{-(N-\nu)/p(\infty)},
\end{aligned}$$

as required. \square

We know the following result.

LEMMA 3.5 ([6, Theorem 1.5]). *Suppose that $p(\cdot)$ satisfies (P1), (P2) and (P3). Then there exists a constant $c_0 > 0$ such that*

$$\|Mf\|_{L^{p(\cdot)}(\mathbf{R}^N)} \leq c_0 \|f\|_{L^{p(\cdot)}(\mathbf{R}^N)}$$

for all $f \in L^{p(\cdot)}(\mathbf{R}^N)$.

Now we are ready to prove Theorem 3.1.

Proof of Theorem 3.1. Let f be a nonnegative function on \mathbf{R}^N such that $\|f\|_{\mathcal{B}^{p(\cdot),\nu}(\mathbf{R}^N)} \leq 1$. For $R \geq 1$, set

$$f = f\chi_{B(0,2R)} + f\chi_{\mathbf{R}^N \setminus B(0,2R)} = f_1 + f_2.$$

First we find from Lemmas 3.2 and 3.4

$$\begin{aligned}
\|Mf_2\|_{L^{p(\cdot)}(B(0,R))} &\leq CR^{-(N-\nu)/p(\infty)} \|1\|_{L^{p(\cdot)}(B(0,R))} \\
&\leq CR^{-(N-\nu)/p(\infty)} R^{N/p(\infty)} = CR^{\nu/p(\infty)}.
\end{aligned}$$

Next we obtain by Lemma 3.5

$$\begin{aligned}
\|Mf\|_{L^{p(\cdot)}(B(0,R))} &\leq \|Mf_1\|_{L^{p(\cdot)}(B(0,R))} + \|Mf_2\|_{L^{p(\cdot)}(B(0,R))} \\
&\leq C \left\{ \|f\|_{L^{p(\cdot)}(B(0,2R))} + R^{\nu/p(\infty)} \right\} \\
&\leq C \left\{ (2R)^{\nu/p(\infty)} + R^{\nu/p(\infty)} \right\} \leq CR^{\nu/p(\infty)},
\end{aligned}$$

so that

$$\sup_{R \geq 1} R^{-\nu/p(\infty)} \|Mf\|_{L^{p(\cdot)}(B(0,R))} \leq C.$$

Thus we establish the required result. \square

REMARK 3.6. If $\nu > N$, then, as in the proof of Theorem 3.1, we find

$$\sup_{R \geq 1} R^{-\nu/p(\infty)} \|M(f\chi_{B(0,R)})\|_{L^{p(\cdot)}(B(0,R))} \leq C.$$

4 Sobolev's inequality

For $\nu \geq 0$, take the integer $k \geq 0$ such that

$$k - 1 \leq \alpha - (N - \nu)/p(\infty) < k \quad (4.1)$$

and consider the generalized Riesz potential

$$I_{\alpha,k}f(x) = \int_{\mathbf{R}^N \setminus B(0,1)} \left\{ I_{\alpha}(x-y) - \sum_{\{\mu:|\mu|\leq k-1\}} \frac{x^{\mu}}{\mu!} (D^{\mu}I_{\alpha})(-y) \right\} f(y) dy$$

for a locally integrable function f on \mathbf{R}^N .

The following estimates are fundamental (see [17] and [19]).

LEMMA 4.1. *Let $k \geq 1$ be an integer.*

(1) *If $2|x| < |y|$, then*

$$\left| I_{\alpha}(x-y) - \sum_{\{\mu:|\mu|\leq k-1\}} \frac{x^{\mu}}{\mu!} (D^{\mu}I_{\alpha})(-y) \right| \leq C|x|^k|y|^{\alpha-N-k};$$

(2) *If $|x|/2 \leq |y| \leq 2|x|$, then*

$$\left| I_{\alpha}(x-y) - \sum_{\{\mu:|\mu|\leq k-1\}} \frac{x^{\mu}}{\mu!} (D^{\mu}I_{\alpha})(-y) \right| \leq C|x-y|^{\alpha-N};$$

(3) *If $1 \leq |y| \leq |x|/2$, then*

$$\left| I_{\alpha}(x-y) - \sum_{\{\mu:|\mu|\leq k-1\}} \frac{x^{\mu}}{\mu!} (D^{\mu}I_{\alpha})(-y) \right| \leq C|x|^{k-1}|y|^{\alpha-N-(k-1)}.$$

LEMMA 4.2. *Let k be the integer defined by (4.1). Then there exists a constant $C > 0$ such that*

$$|I_{\alpha,k}(f\chi_{\mathbf{R}^N \setminus B(0,2R)})(x)| \leq CR^{\alpha-(N-\nu)/p(\infty)}$$

for all $x \in B(0, R)$ with $R \geq 1$ and $f \geq 0$ with $\|f\|_{\mathcal{B}^{p(\cdot),\nu}(\mathbf{R}^N)} \leq 1$.

Proof. Let f be a nonnegative function on \mathbf{R}^N such that $\|f\|_{\mathcal{B}^{p(\cdot),\nu}(\mathbf{R}^N)} \leq 1$. Let $R \geq 1$ and $x \in B(0, R)$. First note from Lemma 4.1 (1) that

$$|I_{\alpha,k}(f\chi_{\mathbf{R}^N \setminus B(0,2R)})(x)| \leq CR^k \int_{\mathbf{R}^N \setminus B(0,2R)} |y|^{\alpha-N-k} f(y) dy.$$

Hence, we have by Lemma 3.3

$$\begin{aligned}
|I_{\alpha,k}(f\chi_{\mathbf{R}^N \setminus B(0,2R)})(x)| &\leq CR^k \sum_{j=1}^{\infty} \int_{B(0,2^{j+1}R) \setminus B(0,2^jR)} |y|^{\alpha-N-k} f(y) dy \\
&\leq CR^k \sum_{j=1}^{\infty} (2^j R)^{\alpha-k} \frac{1}{|B(0,2^{j+1}R)|} \int_{B(0,2^{j+1}R) \setminus B(0,2^jR)} f(y) dy \\
&\leq CR^k \sum_{j=1}^{\infty} (2^j R)^{\alpha-k-(N-\nu)/p(\infty)} \\
&= CR^{\alpha-(N-\nu)/p(\infty)} \sum_{j=1}^{\infty} 2^{j\{\alpha-k-(N-\nu)/p(\infty)\}} \\
&\leq CR^{\alpha-(N-\nu)/p(\infty)},
\end{aligned}$$

as required. \square

LEMMA 4.3. *Let $k \geq 1$ be an integer. Then there exists a constant $C > 0$ such that*

(1) *in case $k-1 < \alpha - (N-\nu)/p(\infty) < k$,*

$$|x|^{k-1} \int_{B(0,|x|/2) \setminus B(0,1)} |y|^{\alpha-N-(k-1)} f(y) dy \leq CR^{\alpha-(N-\nu)/p(\infty)};$$

(2) *in case $k-1 = \alpha - (N-\nu)/p(\infty)$,*

$$|x|^{k-1} \int_{B(0,|x|/2) \setminus B(0,1)} |y|^{\alpha-N-(k-1)} f(y) dy \leq CR^{\alpha-(N-\nu)/p(\infty)} \log R$$

for all $x \in B(0, R)$ with $R \geq 2$ and $f \geq 0$ with $\|f\|_{\mathcal{B}^{p(\cdot),\nu}(\mathbf{R}^N)} \leq 1$.

Proof. Let f be a nonnegative function on \mathbf{R}^N such that $\|f\|_{\mathcal{B}^{p(\cdot),\nu}(\mathbf{R}^N)} \leq 1$. Let $R \geq 2, k \geq 1$ and $x \in B(0, R)$. We may assume that $|x| \geq 2$. We take an integer $j_0 \geq 1$ such that $2^{-j_0-1}|x| < 1 \leq 2^{-j_0}|x|$.

First we show the case $k-1 < \alpha - (N-\nu)/p(\infty) < k$. Then we have by Lemma 3.3

$$\begin{aligned}
&|x|^{k-1} \int_{B(0,|x|/2) \setminus B(0,1)} |y|^{\alpha-N-(k-1)} f(y) dy \\
&\leq |x|^{k-1} \sum_{j=1}^{j_0} \int_{B(0,2^{-j}|x|) \setminus B(0,2^{-j-1}|x|)} |y|^{\alpha-N-(k-1)} f(y) dy \\
&\leq C|x|^{k-1} \sum_{j=1}^{\infty} (2^{-j}|x|)^{\alpha-(k-1)} \frac{1}{|B(0,2^{-j}|x|)|} \int_{B(0,2^{-j}|x|) \setminus B(0,2^{-j-1}|x|)} f(y) dy \\
&\leq CR^{k-1} \sum_{j=1}^{j_0} (2^{-j}R)^{\alpha-(k-1)-(N-\nu)/p(\infty)} \\
&\leq CR^{\alpha-(N-\nu)/p(\infty)}.
\end{aligned}$$

Next we deal with the case $k-1 = \alpha - (N-\nu)/p(\infty)$. Since $j_0 \leq \log|x|/\log 2 < j_0 + 1$, we see from Lemma 3.3 that

$$\begin{aligned} |x|^{k-1} \int_{B(0,|x|/2) \setminus B(0,1)} |y|^{\alpha-N-(k-1)} f(y) dy &\leq CR^{k-1} \sum_{j=1}^{j_0} (2^{-j}R)^{\alpha-(k-1)-(N-\nu)/p(\infty)} \\ &\leq CR^{\alpha-(N-\nu)/p(\infty)} j_0 \\ &\leq CR^{\alpha-(N-\nu)/p(\infty)} \log R, \end{aligned}$$

as required. \square

Set

$$1/p^\sharp(x) = 1/p(x) - \alpha/N.$$

LEMMA 4.4 ([18, Theorem 4.1]). *Suppose $1/p^+ - \alpha/N > 0$. Then there exists a constant $c_1 > 0$ such that*

$$\|I_\alpha f\|_{L^{p^\sharp(\cdot)}(\mathbf{R}^N)} \leq C \|f\|_{L^{p(\cdot)}(\mathbf{R}^N)}$$

for all $f \in L^{p(\cdot)}(\mathbf{R}^N)$ with compact support.

Now we show the Sobolev type inequality for generalized Riesz potentials in the central Morrey spaces of variable exponents, as an extension of Fu, Lin and Lu [9] in the constant exponent case.

THEOREM 4.5 (cf. [9, Proposition 1.1]). *Suppose $1/p^+ - \alpha/N > 0$ and $k-1 < \alpha - (N-\nu)/p(\infty) < k$. Then there exists a constant $C > 0$ such that*

$$\sup_{R \geq 1} R^{-\nu/p(\infty)} \|I_{\alpha,k} f\|_{L^{p^\sharp(\cdot)}(B(0,R))} \leq C$$

for all $f \geq 0$ with $\|f\|_{\mathcal{B}^{p(\cdot),\nu}(\mathbf{R}^N)} \leq 1$.

Proof. Let f be a nonnegative function on \mathbf{R}^N such that $\|f\|_{\mathcal{B}^{p(\cdot),\nu}(\mathbf{R}^N)} \leq 1$. For $R \geq 1$, set

$$f = f\chi_{B(0,2R)} + f\chi_{\mathbf{R}^N \setminus B(0,2R)} = f_1 + f_2.$$

First we find by Lemmas 3.2 and 4.2

$$\begin{aligned} \|I_{\alpha,k} f_2\|_{L^{p^\sharp(\cdot)}(B(0,R))} &\leq CR^{\alpha-(N-\nu)/p(\infty)} \|1\|_{L^{p^\sharp(\cdot)}(B(0,R))} \\ &\leq CR^{\alpha-(N-\nu)/p(\infty)} R^{N/p^\sharp(\infty)} \\ &= CR^{\nu/p(\infty)}. \end{aligned}$$

Next, we see from Lemmas 4.1 and 4.3 (1) that

$$\begin{aligned} &|I_{\alpha,k} f_1(x)| \\ &\leq |I_{\alpha,k}(f\chi_{B(0,2R) \setminus B(0,2|x|)})(x)| + |I_{\alpha,k}(f\chi_{B(0,2|x|) \setminus B(0,|x|/2)})(x)| + |I_{\alpha,k}(f\chi_{B(0,|x|/2) \setminus B(0,1)})(x)| \\ &\leq C \{I_\alpha f_1(x) + R^{\alpha-(N-\nu)/p(\infty)}\} \end{aligned}$$

for $x \in B(0, R)$ since $|x|^k |y|^{\alpha-N-k} \leq C|x-y|^{\alpha-N}$ for $2|x| < |y|$, so that we have by Lemmas 3.2 and 4.4

$$\begin{aligned} \|I_{\alpha,k} f\|_{L^{p^\sharp(\cdot)}(B(0,R))} &\leq \|I_{\alpha,k} f_1\|_{L^{p^\sharp(\cdot)}(B(0,R))} + \|I_{\alpha,k} f_2\|_{L^{p^\sharp(\cdot)}(B(0,R))} \\ &\leq C \left\{ \|f\|_{L^{p(\cdot)}(B(0,2R))} + R^{\nu/p(\infty)} \right\} \\ &\leq C \left\{ (2R)^{\nu/p(\infty)} + R^{\nu/p(\infty)} \right\} \leq CR^{\nu/p(\infty)}, \end{aligned}$$

so that

$$\sup_{R \geq 1} R^{-\nu/p(\infty)} \|I_{\alpha,k} f\|_{L^{p^\sharp(\cdot)}(B(0,R))} \leq C.$$

Thus we completes the proof. \square

REMARK 4.6. Suppose $1/p^+ - \alpha/N > 0$ and $k - 1 = \alpha - (N - \nu)/p(\infty)$. Then there exists a constant $C > 0$ such that

$$\sup_{R \geq 2} R^{-\nu/p(\infty)} (\log R)^{-1} \|I_{\alpha,k} f\|_{L^{p^\sharp(\cdot)}(B(0,R))} \leq C$$

for all $f \geq 0$ with $\|f\|_{\mathcal{B}^{p(\cdot),\nu}(\mathbf{R}^N)} \leq 1$.

5 Exponential integrability

Our aim in this section is to discuss the exponential integrability.

THEOREM 5.1. *Let $p = N/\alpha$ and $k - 1 < \alpha - (N - \nu)/p < k$. Then there exist constants $c_1, c_2 > 0$ such that*

$$\sup_{R \geq 1} R^{-N} \int_{B(0,R)} \exp(\{c_1 R^{-\nu/p} |I_{\alpha,k} f(x)|\}^{p'}) dx \leq c_2$$

for all $f \geq 0$ with $\|f\|_{\mathcal{B}^{p,\nu}(\mathbf{R}^N)} \leq 1$.

Proof. Let f be a nonnegative function on \mathbf{R}^N such that $\|f\|_{\mathcal{B}^{p,\nu}(\mathbf{R}^N)} \leq 1$ and let $x \in B(0, R)$. For $R \geq 1$, set

$$f = f\chi_{B(0,2R)} + f\chi_{\mathbf{R}^N \setminus B(0,2R)} = f_1 + f_2.$$

For $0 < \delta \leq R$, write

$$\begin{aligned} I_{\alpha} f_1(x) &= \int_{B(x,\delta)} |x-y|^{\alpha-N} f(y) dy + \int_{B(0,2R) \setminus B(x,\delta)} |x-y|^{\alpha-N} f(y) dy \\ &= U_1(x) + U_2(x). \end{aligned}$$

First we find

$$U_1(x) \leq C\delta^{\alpha} M f_1(x).$$

Next we have by Hölder's inequality

$$U_2(x) \leq C(\log(2R/\delta))^{1/p'} \|f_1\|_{L^p(B(0,2R))},$$

so that

$$I_\alpha f_1(x) \leq C \left\{ \delta^\alpha M f_1(x) + (\log(2R/\delta))^{1/p'} R^{\nu/p} \right\}.$$

Here, letting $\delta/(2R) = \{R^{-\nu/p+\alpha} M f_1(x)\}^{-1/\alpha} (\log(R^{-\nu/p+\alpha} M f_1(x)))^{1/(\alpha p')} < 1$, we establish

$$I_\alpha f_1(x) \leq C (\log(R^{-\nu/p+\alpha} M f_1(x)))^{1/p'} R^{\nu/p};$$

if $\{R^{-\nu/p+\alpha} M f_1(x)\}^{-1/\alpha} (\log(R^{-\nu/p+\alpha} M f_1(x)))^{1/(\alpha p')} \geq 1$, then, letting $\delta = R$, we have

$$I_\alpha f_1(x) \leq C R^{\nu/p}.$$

As in the proof of Theorem 4.5, we see from Lemmas 4.1 and 4.3 (1) that

$$|I_{\alpha,k} f_1(x)| \leq C \{I_\alpha f_1(x) + R^{\alpha-(N-\nu)/p}\} = C \{I_\alpha f_1(x) + R^{\nu/p}\}$$

for $x \in B(0, R)$, since $\alpha = N/p$. Therefore, we obtain

$$|I_{\alpha,k} f_1(x)| \leq C \{(\log(e + R^{-\nu/p+\alpha} M f_1(x)))^{1/p'} R^{\nu/p} + R^{\nu/p}\}.$$

On the other hand, we obtain by Lemma 4.2

$$|I_{\alpha,k} f_2(x)| \leq C R^{\alpha-(N-\nu)/p} = C R^{\nu/p},$$

since $\alpha = N/p$. Hence, we find

$$\{c_1 R^{-\nu/p} |I_{\alpha,k} f(x)|\}^{p'} \leq \log(e + R^{(N-\nu)/p} M f_1(x)),$$

so that we have by boundedness of maximal operators on $L^p(\mathbf{R}^N)$

$$\begin{aligned} \int_{B(0,R)} \exp(\{c_1 R^{-\nu/p} |I_{\alpha,k} f(x)|\}^{p'}) dx &\leq C \int_{B(0,R)} [1 + R^{N-\nu} \{M f_1(x)\}^p] dx \\ &\leq C \left(R^N + R^{N-\nu} \int_{\mathbf{R}^N} f_1(y)^p dy \right) \\ &\leq C R^N, \end{aligned}$$

as required. \square

REMARK 5.2. Let $p = N/\alpha$ and $k-1 = \alpha - (N-\nu)/p$. Then there exist constants $c_1, c_2 > 0$ such that

$$\sup_{R \geq 2} R^{-N} \int_{B(0,R)} \exp(\{c_1 R^{-\nu/p} (\log R)^{-1} |I_{\alpha,k} f(x)|\}^{p'}) dx \leq c_2$$

for all $f \geq 0$ with $\|f\|_{\mathcal{B}^{p,\nu}(\mathbf{R}^N)} \leq 1$.

REMARK 5.3. If $p^- \geq p(\infty)$, then $\mathcal{B}^{p(\cdot),\nu}(\mathbf{R}^N) \subset \mathcal{B}^{p(\infty),\nu}(\mathbf{R}^N)$, and moreover

$$\|f\|_{\mathcal{B}^{p(\infty),\nu}(\mathbf{R}^N)} \leq C \|f\|_{\mathcal{B}^{p(\cdot),\nu}(\mathbf{R}^N)}.$$

In fact, for $R \geq 1$ and $a > N/p(\infty)$,

$$\begin{aligned} R^{-\nu} \int_{B(0,R)} |f(x)|^{p(\infty)} dx &= R^{-\nu} \int_{\{x \in B(0,R): |f(x)| \geq 1\}} |f(x)|^{p(\infty)} dx \\ &\quad + R^{-\nu} \int_{\{x \in B(0,R): (1+|x|)^{-a} < |f(x)| \leq 1\}} |f(x)|^{p(\infty)} dx \\ &\quad + R^{-\nu} \int_{\{x \in B(0,R): |f(x)| \leq (1+|x|)^{-a}\}} |f(x)|^{p(\infty)} dx \\ &\leq R^{-\nu} \int_{\{x \in B(0,R): |f(x)| \geq 1\}} |f(x)|^{p(x)} dx \\ &\quad + R^{-\nu} \int_{\{x \in B(0,R): (1+|x|)^{-a} < |f(x)| \leq 1\}} |f(x)|^{p(x)} |f(x)|^{p(\infty)-p(x)} dx \\ &\quad + CR^{-\nu} \int_{B(0,R)} (1+|x|)^{-ap(\infty)} dx \\ &\leq C \left\{ R^{-\nu} \int_{B(0,R)} |f(x)|^{p(x)} dx + R^{-\nu} \right\} \\ &\leq C \end{aligned}$$

when $\|f\|_{\mathcal{B}^{p(\cdot),\nu}(\mathbf{R}^N)} \leq 1$.

References

- [1] D. R. Adams and L. I. Hedberg, *Function Spaces and Potential Theory*, Springer, 1996.
- [2] A. Almeida and D. Drihem, Maximal, potential and singular type operators on Herz spaces with variable exponents. *J. Math. Anal. Appl.* **394** (2012), no. 2, 781-795.
- [3] J. Alvarez, M. Guzmán-Partida and J. Lakey, Spaces of bounded λ -central mean oscillation, Morrey spaces, and λ -central Carleson measures, *Collect. Math.* **51** (2000), 1-47.
- [4] A. Beurling, Construction and analysis of some convolution algebras, *Ann. Inst. Fourier* **14** (1964), 1-32.
- [5] B. Bojarski and P. Hajłasz, Pointwise inequalities for Sobolev functions and some applications, *Studia Math.* **106**(1) (1993), 77-92.
- [6] D. Cruz-Uribe, A. Fiorenza and C. J. Neugebauer, The maximal function on variable L^p spaces, *Ann. Acad. Sci. Fenn. Math.* **28** (2003), 223-238; *Ann. Acad. Sci. Fenn. Math.* **29** (2004), 247-249.

- [7] L. Diening, P. Harjulehto, P. Hästö and M. Růžička, Lebesgue and Sobolev spaces with variable exponents, Lecture Notes in Mathematics, **2017**, Springer, Heidelberg, 2011.
- [8] H. Feichtinger, An elementary approach to Wiener's third Tauberian theorem on Euclidean n -space, Proceedings, Conference at Cortona 1984, Symposia Mathematica 29 (Academic Press, New York, 1987), 267–301.
- [9] Z. Fu, Y. Lin and S. Lu, λ -central BMO estimates for commutators of singular integral operators with rough kernels, Acta Math. Sin. (Engl. Ser.) **24** (2008), no. 3, 373–386.
- [10] J. García-Cuerva, Hardy spaces and Beurling algebras, J. London Math. Soc. **39** (1989), 499–513.
- [11] J. García-Cuerva and M. J. L. Herrero, A theory of Hardy spaces associated to the Herz spaces, Proc. London Math. Soc. **69** (1994), 605–628
- [12] C. Herz, Lipschitz spaces and Bernstein's theorem on absolutely convergent Fourier transforms, J. Math. Mech. **18** (1968), 283–324.
- [13] Y. Komori, Notes on singular integrals on some inhomogeneous Herz spaces, Taiwanese J. Math. **8** (2004), 547–556.
- [14] J. L. Lewis, On very weak solutions of certain elliptic systems, Comm. Partial Differential Equations **18**(9) (10) (1993), 1515–1537.
- [15] K. Matsuoka and E. Nakai, Fractional integral operators on $B^{p,\lambda}$ with Morrey-Campanato norms, Function Spaces IX (Krakow, Poland, 2009), 249–264, Banach Center Publications, Vol.92, Institute of Mathematics, Polish Academy of Sciences, Warszawa, 2011.
- [16] Y. Mizuta, Potential theory in Euclidean spaces, Gakkōtoshō, Tokyo, 1996.
- [17] Y. Mizuta, Integral representations, differentiability properties and limits at infinity for Beppo Levi functions, Potential Analysis **6** (1997), 237–276.
- [18] Y. Mizuta, E. Nakai, T. Ohno and T. Shimomura, Maximal functions, Riesz potentials and Sobolev embeddings on Musielak-Orlicz-Morrey spaces of variable exponent in \mathbf{R}^n , Rev. Mat. Complut. **25** (2012), no. 2, 413–434.
- [19] T. Shimomura and Y. Mizuta, Taylor expansion of Riesz potentials, Hiroshima Math. J. **25** (1995), 595–621.
- [20] E. M. Stein, Singular integrals and differentiability properties of functions, Princeton Univ. Press, Princeton, 1970.
- [21] N. Wiener, Generalized harmonic analysis, Acta Math. **55** (1930), 117–258.
- [22] N. Wiener, Tauberian theorems, Ann. Math. **33** (1932), 1–100.

*Department of Mechanical Systems Engineering
Hiroshima Institute of Technology
2-1-1 Miyake Saeki-ku Hiroshima 731-5193, Japan
E-mail : yoshihiromizuta3@gmail.com*

and

*Faculty of Education and Welfare Science
Oita University
Dannoharu Oita-city 870-1192, Japan
E-mail : t-ohno@oita-u.ac.jp*

and

*Department of Mathematics
Graduate School of Education
Hiroshima University
Higashi-Hiroshima 739-8524, Japan
E-mail : tshimo@hiroshima-u.ac.jp*